



# Glossary of Mathematical Terms



**Introduction:** At Sylvan, we recognize that mathematics is a language that can be learned. This glossary of mathematical terms and definitions includes several hundred of the most important mathematical terms in use today. This glossary is intended for use by all members of the Sylvan Learning family.



## Glossary of Mathematics Terms

Term	Definition
4-step method for problem solving	A problem-solving method that can be applied to any math problem. Students are taught the steps as: Understand, Plan, Solve, and Look Back.
absolute value	The distance a number is from zero on a number line. The absolute value of $x$ is denoted by $ x $ . Example: $ -5  = 5$ ; $ 5  = 5$ .
acute angle	An angle whose measure is greater than $0^\circ$ and less than $90^\circ$ .
acute triangle	A triangle with three acute angles.
addend	A number being added to obtain a sum.
additive inverse	Also called the opposite of a number. The sum of any number and its additive inverse is zero. Example: the additive inverse of 5 is $-5$ and the additive inverse of $-5$ is 5.
adjacent angles	A pair of angles with a common vertex and exactly one common side.
algebra	A generalization and an extension of the concepts of arithmetic. In algebra, variables are used to represent numbers and quantities.
algebraic equation	A mathematical sentence that includes one or more variables and contains two expressions joined by an equal sign.
algebraic expression	A variable or a combination of one or more variables and numbers with one or more operations. Examples: $x$ , $x + 9$ , $2x - 3$ , and $4x^2 - 3y^2$ .
algorithm	A set of step-by-step directions for a procedure. Examples: Partial Sums algorithm, Column Addition Algorithm, Trade First Subtraction algorithm, etc.
altitude	A segment that extends from a vertex of a figure and meets a base at a right angle. Also called the height.
angle	A figure formed by two rays with a common endpoint. The common endpoint is called the vertex of the angle.
arc	A part of a circle consisting of two endpoints and all of the points on the circle between the two endpoints.
area	The number of square units needed to cover a region. $A = \pi r^2$
associative property	The grouping of addends does not change the sum. $(a + b) + c = a + (b + c)$ The grouping of factors does not change the product. $(ab)c = a(bc)$
axes	The vertical and horizontal lines of reference that divide the coordinate plane into quadrants. The horizontal axis is the $x$ -axis, and the vertical axis is the $y$ -axis.
bar graph	A graph that uses vertical or horizontal bars to represent relationships among data.
binomial	A polynomial that consists of two terms, or two monomials.

Term	Definition
box-and-whisker plot	A graphical method for showing the median, quartiles, and extremes of data.
Cartesian coordinate system	The system of locating points on a plane formed by a horizontal line ( $x$ -axis) intersecting a vertical line ( $y$ -axis). The plane is divided into four quadrants in which ordered pairs, or points, are identified and plotted by their distance from each axis.
central angle	An angle that has its vertex at the center of a circle. The measure of the central angle equals the measure of the minor arc.
certain event	An event that must happen. It has a probability of 1.
chart	A display used to organize information or data.
chord	A line segment whose endpoints lie on a circle.
circle	The set of all points in a plane that are a given distance (radius) from a fixed point (center).
circle graph	A type of graph where a circle represents the total amount and segments denote the data. It is also known as a pie chart.
circumference	The distance around a circle. $C = \pi d$
coefficient	A constant used as a factor in an algebraic expression. In the expression $2x + 3y$ , the coefficient of $x$ is 2, and the coefficient of $y$ is 3.
coincident	Two or more geometric figures that share all points. Example: two coincident lines would have equations that are equivalent.
combination	An arrangement of elements or events in which order does not matter.
common factor	A common factor of two natural numbers is a number that is a factor of both numbers. Example: 7 is a common factor of 35 and 56, because $35 = 7 \times 5$ and $56 = 7 \times 8$
commutative property	The order of addends does not change the sum. $a + b = b + a$ The order of factors does not change the product. $ab = ba$
compatible numbers	Pairs of numbers that can easily be added, subtracted, multiplied, or divided using mental math.
complementary angles	A pair of angles the sum of whose measures is $90^\circ$ .
composite figure	A figure made up of two or more basic two-dimensional objects.
composite number	An integer greater than 1 with more than two factors.
concave polygon	A polygon that has at least one diagonal outside the polygon.
congruent	Having the same shape and the same size.
congruent angles	Two or more angles that have the same degree measure.
consecutive numbers	Whole numbers that follow in order. Example: 3, 4, and 5
consecutive sides of a polygon	Adjacent sides of a polygon.
convex polygon	A polygon that has all of its diagonals inside the polygon.

Term	Definition
coordinate plane or system	A plane formed by a horizontal line ( $x$ -axis) that intersects a vertical line ( $y$ -axis). The plane is divided into four quadrants in which ordered pairs or points can be identified and plotted.
coordinates	The ordered pair $(x, y)$ used to represent a point on the coordinate plane.
correlation	A relationship between two sets of data. A positive correlation exists when one set increases or decreases as the other set does the same. A negative correlation exists when one set increases or decreases as the other set does the opposite.
corresponding angles in parallel lines cut by a transversal	When parallel lines are cut by a transversal, angles in the same position relative to the parallel lines and transversal are corresponding and congruent.
corresponding angles of similar triangles	In similar triangles, the angles in the same relative location of the two triangles are corresponding and congruent. If $\triangle ABC \sim \triangle XYZ$ , then $\angle A \approx \angle X$ and $m\angle A = m\angle X$ .
cube	A rectangular prism with all edges and faces congruent.
cubed	Another way to express a number or variable raised to the third power.
cubic units	The units used to measure the space inside a three-dimensional figure.
cylinder	A solid with congruent and parallel circular bases.
data	Information about a situation, group, or event.
data set	A collection of facts or experimental results.
decagon	A 10-sided polygon.
decimal	A number written in standard notation, usually containing a decimal point. Example: 5.25 and 0.36 are decimal numbers.
denominator	The number of equal parts or groups into which the whole or group is divided. The denominator can be any number except zero. Example: In the fraction $\frac{a}{b}$ , $b$ is the denominator.
diagonal	A line segment that connects two non-adjacent vertices in a polygon.
diameter	A line segment passing through the center of a circle and connecting two points on the circle. The diameter is equal to twice the length of the radius.
difference	The result of subtracting two numbers. Example: $7 - 5 = 2$ , the difference is 2.
digit	Any of the numerals 0 through 9.
dilation	A transformation that enlarges or reduces a figure.
dimensional analysis	The process of using unit factors to convert measurements.
direct measure	Measuring an object by physical means. This can be done with a device such as a ruler.
direct variation	The relationship between two variables such that their ratio remains constant.

Term	Definition
distance	The measurement between two points in a plane.
distance formula	The distance traveled ( $d$ ) in a given time ( $t$ ) at a rate ( $r$ ) is given by the formula $d = rt$ .
distributive property	The product of a factor and a sum is equal to the sum of the products. Example: $a(b + c) = ab + ac$ .
dividend	The amount that is divided into parts. Example: Given $a \div b = c$ or $b \overline{)c}$ , $a$ is the dividend.
divisible	An integer is divisible by another non-zero integer if the division leaves a remainder of zero.
division	The inverse operation of multiplication. Division is performed on two numbers to obtain a third number, called the quotient. Example: $6 \div 2 = 3$ , the number 6 is the dividend, the number 2 is the divisor, and the number 3 is the quotient.
division law of exponents	To divide two powers that have the same base, subtract the exponents. Example: $\frac{x^5}{x^3} = x^2$
divisor	The number by which a dividend is divided. Example: Given $a \div b = c$ or $b \overline{)a}$ , $b$ is the divisor.
domain	The set of all possible values for the unknown in an open sentence or a function.
dotplot	A graphical display in which the frequency of each result is indicated by the number of dots above that result.
double bar graph	A bar graph used to compare two sets of data.
double line graph	A graph with two lines representing different sets of data.
edge	The line along which two faces of a solid figure meet.
edge of a graph	A line segment that connects two vertices in a graph.
element	An item in a set.
endpoints	The initial point and end point of a line segment.
equal	The term used to describe a relationship between two numbers or expressions that have exactly the same value.
equation	A mathematical sentence that uses an equal sign to show that two quantities are equal.
equilateral triangle	A triangle with three sides of equal length and three angles of equal measure. Each angle in an equilateral triangle has a measure of $60^\circ$ .
equivalent	Equal in value, but in a different form. Example: $\frac{3}{4}$ , 0.75, and 75% are equivalent.

Term	Definition
equivalent fractions	Fractions that name the same number. Example: $\frac{3}{4}$ and $\frac{6}{8}$ are equivalent fractions.
equivalent ratios	Ratios that make the same comparison. Example: 5 to 10 and 1 to 2 are equivalent ratios.
estimate	An approximation, or rough calculation, of a quantity or measurement.
even integer	An integer that can be written as the sum of two equal integers. An even number can be expressed in the form $2x$ , where $x$ is any integer. Even integers end in 0, 2, 4, 6, and 8.
even number	A whole number that is divisible by 2. Example: 0, 2, 4, 6, ...
expanded form	The expanded form of an algebraic expression is the equivalent expression without parentheses. Example: the expanded form of $(a + b)^2$ is $a^2 + 2ab + b^2$ .
expanded notation	A way of writing a number to show the value of each digit. Example: $34,718 = 30,000 + 4,000 + 700 + 10 + 8$ .
expenses	The total amount of money spent, including all costs.
experimental probability	Probability of an event based on the statistical results of an actual experiment that has already been performed. Also called empirical probability.
exponent	A symbol written above and to the right of a base that tells how many times the base is used as a factor. Example: $5^3$ , the exponent is 3.
exponential function	A function commonly used to study growth and decay. It has the form $f(x) = a^x$ , when $a > 0$ .
expression	A mathematical arrangement of numbers and variables connected by operations such as addition, subtraction, multiplication, and division. An expression does not contain an equal sign.
exterior angle of a polygon	An angle on the outside of a polygon formed by one side of the polygon and the extension of a second side of the polygon.
face	A flat side of a solid figure.
factor	A number that divides exactly into another number with no remainder. When two or more factors are multiplied, they form a product. Example: $3 \times 6 = 18$ , 3 and 6 are factors, and 18 is the product.
factor	To write a number or expression as a product of its factors.
factor tree	A graphical tool used to help determine the prime factors of a number.
factored form	A way to write numbers using factors. Example: $2 \times 3 \times 5$ is the factored form of 30.

<b>Term</b>	<b>Definition</b>
factorial	The product of all of the positive integers less than or equal to a number. The symbol for factorial is $!$ . Example: $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$
formal mathematical language	A written language that relies on precise mathematical terms and definitions to communicate ideas.
fraction	A number in the form $\frac{a}{b}$ , where $a$ and $b$ are whole numbers and $b$ is not 0. Fractions are used to name parts of a whole object or part of a whole collection of objects, or to compare two quantities.
fractional equation	An equation that includes fractional coefficients. Example: $\frac{a}{4} + \frac{a}{3} = 5$
function	A mathematical relationship between two sets in which each element of the domain is matched with exactly one element of the range.
function table	Pairs of $x$ and $y$ values arranged in a table to represent a function, relationship, sequence, or pattern.
fundamental counting principle	If one event has $p$ possible outcomes and an independent event has $q$ possible outcomes, then the first event followed by the second event has $p \times q$ possible outcomes.
geoboard	A board with pegs aligned in grid fashion that permits rubber bands to be wrapped around pegs to form geometric figures.
geometric sequence	A sequence of numbers that is generated by multiplying each new term by the same number. Example: In the sequence $\{1, 4, 16, 64, 256, \dots\}$ each term has been multiplied by 4 to generate the next term in the sequence.
geometry	The study of the properties, measurement, and relationships of points, lines, angles, planes, and solids.
glyph	A pictorial form of data collection used to organize and analyze data.
graph	A data display that uses ordered pairs, bars, lines, circles, or pictures to show a data pattern or as a model to predict future values.
graphing calculator	A calculator that will store and draw the graphs of several functions at once.
greater than	The symbol used to denote that one quantity is greater than another. Example: $8 > 5$
greatest common factor (GCF)	The largest number that is a factor of two or more numbers. Example: the GCF of 36 and 24 is 12, since $36 = 12 \times 3$ and $24 = 12 \times 2$ .
guess and check	A problem-solving strategy that involves testing possible answers and using the results of those tests to determine the answer.

Term	Definition
height of a prism or cylinder	The perpendicular distance between the bases.
height of a pyramid or cone	The length of the perpendicular segment from the vertex to the base.
height of a trapezoid	The perpendicular distance between the bases.
height of a triangle	The perpendicular segment from a vertex to the base. Also called the altitude.
histogram	A type of graph that uses adjacent vertical bars to show frequencies of data.
horizontal axis	The axis running left to right, often denoted as the $x$ -axis in a coordinate plane.
hundredth	Each of one hundred equal parts.
hypotenuse	The side opposite the right angle in a right triangle. The hypotenuse is the longest side of a right triangle.
identity	A mathematical equation that is true for all real numbers. Example: $x + x = 2x$
identity property of addition	The sum of any number and 0 is that number. For all real numbers $a$ , $a + 0 = 0 + a = a$ . Zero is called the additive identity.
identity property of multiplication	The product of any real number and 1 is that number. The real number 1 is the multiplicative identity. $1 \times a = a \times 1 = a$
image	The result of a transformation.
impossible event	An event that cannot happen, it has a probability of 0.
independent events	Two or more events in which the outcome of any one event does not affect the outcome of any of the other events.
independent variable	A mathematical term or variable whose value is given or defined first, and determines the value of the dependent variable. In a function, the domain lists the possible values of the independent variable.
indirect measure	Measuring an object on the basis of another object whose measure is known.
inequality	A mathematical sentence that uses symbols such as for greater than ( $>$ ) and less than ( $<$ ) to compare values.
inscribed angle	An angle formed by two chords of a circle with a common endpoint. The measure of the inscribed angle is half the measure of the intercepted arc.
inscribed polygon	A polygon whose sides are all chords of a circle.
integers	The set of whole numbers and their opposites. Zero is an integer, and is neither positive nor negative. $\{\dots -3, -2, -1, 0, 1, 2, 3, \dots\}$
intercepted arc	An arc that is formed by an inscribed angle. The arc falls in the interior of the angle, with endpoints on the rays of the inscribed angle.

Term	Definition
interest	A sum of money paid when borrowing or saving money. Simple interest is calculated by multiplying the principle (P) by the rate (r) and the amount of time (t). $I = Prt$
interior angles	Angles on the inside of a polygon formed by two consecutive sides of the polygon.
intersection	The point where two lines cross.
interval	The constant space between grid lines on the axes of a graph.
inverse operations	Operations that are opposites. Addition is the inverse operation of subtraction, and multiplication is the inverse operation of division.
inverse property of addition	The sum of any real number and its opposite is zero. The real number $-a$ is the additive inverse of $a$ . $a + (-a) = (-a) + a = 0$
inverse property of multiplication	The product of any nonzero real number and its reciprocal is 1. The real number $\frac{1}{a}$ is the multiplicative inverse of the nonzero real number $a$ . $a\left(\frac{1}{a}\right) = \left(\frac{1}{a}\right)a = 1$
inverse variation	The relationship between two variables such that the ratio of one variable to the reciprocal of the other is a constant.
isosceles right triangle	A right triangle whose two legs have equal lengths. The angles of an isosceles right triangle measure $45^\circ$ , $45^\circ$ , and $90^\circ$ .
isosceles trapezoid	A trapezoid with non-parallel sides congruent.
isosceles triangle	A triangle with at least two congruent sides and angles.
kite	A four-sided polygon with exactly two distinct pairs of congruent consecutive sides.
lateral area	The sum of the areas of the surfaces, excluding the bases, of a three-dimensional figure.
lateral edge	A line formed by the intersection of two lateral faces of a prism.
lateral faces	The faces of a prism that are not bases.
least common multiple (LCM)	The smallest number that is a multiple of two or more numbers. Example: LCM of 15 and 20 is 60.
legend	A key that explains the symbols used in a graph.
legs	In a right triangle, the two sides that intersect to form the right angle.
length	A dimension of a rectangle, usually the longer side.
less than	The symbol used to denote that one quantity is less than another. Example: $5 < 8$
like terms	Terms with the same variables and the same powers of those variables. Example: $2x^3$ and $-5x^3$ are like terms.
likely	A result that will probably, but not definitely, occur.

<b>Term</b>	<b>Definition</b>
line	A set of points that form a straight path extending infinitely in both directions.
line graph	A graph with a horizontal and vertical axis that represents data as a continuous line or curve.
line segment	The set of points between two given points on a line.
linear algebra	The branch of mathematics that considers vectors, systems of linear equations, and linear transformations.
linear equation	An equation that represents a straight line.
linear function	A function whose general equation is $f(x) = mx + b$ , where $m$ and $b$ are constants.
linear measurement	Measurement in a straight line.
linear pair of angles	A pair of adjacent angles whose distinct sides lie on a straight line. The sum of the measures of a linear pair of angles is $180^\circ$ .
liter	The basic unit for measuring capacity within the metric system. One liter equals 1,000 milliliters.
magic square	A square containing integers arranged in an equal number of rows and columns so that each row, column, and diagonal has the same sum.
major arc	An arc whose measure is more than the measure of a semicircle.
manipulatives	Materials that allow students to explore mathematical concepts in a concrete mode.
mass	The amount of matter that an object contains.
mathematical model	A representation in the mathematical world of some phenomenon in the real world.
maximum	The largest value in a data set.
maximum point	The highest point on a graph.
mean	The average of a set of numbers. The mean can be found by taking the sum of the numbers in a set of data divided by the number of pieces of data.
measures of central tendency	Values that summarize a set of numerical data. The most common measures of central tendency are mean, median, mode, and range.
median	The middle number in a set of numbers that are arranged from least to greatest. The median of a set with an even number of elements is the arithmetic mean of the two middle elements.
mental computation	A calculation done without pencil, paper, calculator, or computer.
meter	A unit of length in the metric system. One meter equals 100 centimeters.
metric system	A system of measurement that is based on units of 10. The basic units of the metric system are the meter, liter, and gram.
midpoint	The point on a line segment that is equidistant from the endpoints.

Term	Definition
minimum	The smallest value in a data set.
minimum point	The lowest point on a graph.
minor arc	An arc whose measure is less than $180^\circ$ .
minuend	The first number in an expression of subtraction. Example: In the expression $5 - 3 = 2$ , the minuend is 5.
mixed number	A number consisting of a whole number and a fraction Example: $3\frac{1}{2}$
mode	In statistics, the value that occurs most frequently in a given set of numbers.
monomial	A single term that is made up of a number, a variable, or the product of numbers and variables. Example: 4, $2x$ , and $-5x^2y^3$ are monomials.
multiple	A multiple of a number is the product of that number and any nonzero whole number.
multiplication	An operation on two numbers, called factors, to obtain a third number called the product. Example: In the equation $5 \times 2 = 10$ , the numbers 5 and 2 are factors, and 10 is the product.
multiplication laws of exponents	To multiply two powers that have the same base, add the exponents. Example: $x^3 \cdot x^5 = x^8$
multiplicative inverse	The opposite of a number in a multiplication operation. The product of any number and its multiplicative inverse is one. The multiplicative inverse is also called the reciprocal. Example: the multiplicative inverse of $\frac{2}{3}$ is $\frac{3}{2}$ .
natural numbers	The set of positive integers, not including zero. Natural numbers are also known as counting numbers. $\{1, 2, 3, 4, 5, \dots\}$
negative integer	An integer that is less than zero. $\{\dots -5, -4, -3, -2, -1\}$
net of a polyhedron	A two-dimensional diagram of a polyhedron.
nonagon	A nine-sided polygon.
nonstandard unit	Unit of measurement expressed in terms of objects, such as paper clips, sticks of gum, shoes, etc.
notation	Symbols and characters used to express values, operations, and relationships in mathematical expressions.
number line	A line on which points are marked off at regular intervals and labeled with ordered numbers.
numerator	When a whole is divided into a number of equal parts, the number of equal parts being considered is the numerator. Example: In the fraction $\frac{a}{b}$ , $a$ is the numerator.
numerical expression	A mathematical phrase that contains numbers and operation symbols, but no variables.
obtuse angle	An angle whose measure is between $90^\circ$ and $180^\circ$ .

Term	Definition
obtuse triangle	A triangle with one angle greater than $90^\circ$ .
odd number	A whole number that is not divisible by 2.
odds	The ratio of the number of favorable outcomes to the number of unfavorable outcomes.
one-to-one correspondence	When each element of a set is assigned to one element of a second set, and each element of the second set is assigned to one element of the first set, the mapping is called a one-to-one correspondence.
open sentence	A statement that contains at least one unknown. It becomes true or false when a quantity is substituted for the unknown. Example: $2x + 3 = 12$ or $5 + \underline{\quad} = 9$
operation	A process performed on numbers and expressions. The basic operations are addition, subtraction, multiplication, and division.
opposite integers	Two integers that are the same distance from zero on a number line, in opposite directions.
order of operations	A rule indicating the order in which operations should be performed when there are several operations in an expression. Perform operations inside grouping symbols first, and then simplify exponents. Next perform multiplication and division from left to right, and finally perform addition and subtraction from left to right. The acronym PEMDOS refers to the specified order.
ordered pair	An ordered pair of numbers $(x, y)$ used to name a point on a coordinate plane.
origin	The point $(0, 0)$ on the coordinate plane at the intersection of the $x$ -axis and the $y$ -axis.
outcome	The result of an experiment.
parabola	A curve in which every point is the same distance from a fixed point (focus) and a fixed line (directrix). The equation of a parabola can be expressed as $y = a(x - h)^2 + k$ , where $(h, k)$ is the vertex of the parabola. The graph is concave up if $a > 0$ and concave down if $a < 0$ .
parallel lines	Lines in the same plane and never intersect.
parallelogram	A four-sided polygon with opposite sides parallel and congruent.
patterns	Regularities in situations such as those in nature, events, shapes, designs, and sets of numbers. Example: spirals on a pineapple, geometric designs in quilts, the number sequence 2, 4, 6, 8, ...
percent	A ratio where the comparison is made out of 100. Example: $17\% = \frac{17}{100}$

Term	Definition
percent decrease	The percent represented by the ratio of the amount of decrease to the original amount.
percent increase	The percent represented by the ratio of the amount of increase to the original amount.
perfect square	A whole number that can be renamed as the square of an integer. Example: 16 is a perfect square because $16 = 4^2$
perfect square trinomial	A trinomial in the form $a^2 + 2ab + b^2$ , which can be factored as $(a + b)^2$ .
perimeter	The distance around a polygon. This distance can be found by adding the lengths of the sides.
perpendicular bisector of a line segment	A line, ray, or line segment that is perpendicular to a given line segment, and divides the line segment into two equal parts.
perpendicular lines	Lines that intersect to form right angles.
pi	The ratio of the circumference of a circle to its diameter. The symbol for pi is $\pi$ . It can be approximated as 3.14 or $\frac{22}{7}$ .
pictograph	A type of graph in which pictures are used to represent values.
pie chart	A type of graph where a circle represents the total amount and segments denote the data. It is also known as a circle graph.
pint	A dry or liquid measure of capacity. Two cups equal 1 pint and two pints equal one quart.
place value	The value of a digit depending on its position in a numeral. In the numeral 4695, the 6 is in the hundreds place, which gives it a value of 600.
plane	A two-dimensional geometric object that extends without boundary.
plane figure	A figure that lies in only one plane. Circles and polygons are plane figures.
plotting points	Locating points, lines, or curves by means of placing coordinates on a grid.
point	A geometric object with position but without length, width, or depth.
polygon	A many-sided closed figure in which each side is a line segment.
polyhedron	A three-dimensional closed figure whose faces are polygons.
polynomial	In algebra, a sum of one or more monomials. Example: $-2x$ , $3x^2 + 5x - 7$ or $a^2b - ab^2$ .
positive integer	An integer that is greater than zero. $\{1, 2, 3, \dots\}$
power	Another name for exponent.
prediction	A guess about the outcome or result of an event or action based on the available information.

Term	Definition
prime factorization	The expression of a composite number as the product of its prime factors. Example: $2^3 \cdot 3 \cdot 5^2$ is the prime factorization of 600.
prime number	A whole number greater than one that has exactly two factors. Example: The first five prime numbers are 2, 3, 5, 7, and 11.
principal	A sum of money that increases by a certain rate of interest.
principal square root	The positive square root of a number. The notation $\sqrt{36}$ indicates the principal square root, 6.
prism	A solid formed by three or more lateral faces that intersect with bases.
probability	The likelihood or chance of an event occurring. Probability is expressed as a number from 0 to 1. A probability of 0 means that the event cannot occur, while a probability of 1 means that the event is certain to occur.
probability formula for geometric regions	The probability that a random point is located in a particular part, or subregion, of a larger region is given by the formula $P(E) = \frac{\text{measure of region in the event}}{\text{measure of entire region}}.$
problem solving	Finding ways to reach a goal when no routine path is apparent.
product	The result of multiplying two or more numbers.
property of proportions	In the proportion $\frac{a}{b} = \frac{c}{d}$ , the product of the means $a \cdot d$ is equal to the product of the extremes $b \cdot c$ . This can be expressed as $a \cdot d = b \cdot c$ .
proportion	An equation stating that two ratios are equivalent.
protractor	A tool used to measure angles in degrees.
pyramid	A solid formed by the intersection of a polygonal base with triangular faces that meet at a vertex.
Pythagorean theorem	The sum of the squares of the lengths of the legs in a right triangle is equal to the square of the length of the hypotenuse. Example: In a right triangle with legs of length $a$ and $b$ and $c$ as the length of the hypotenuse $a^2 + b^2 = c^2$ .
quadrant	One of the four sections or quarters of a coordinate plane.
quadratic equation	An equation of degree 2 that has at most two solutions. Quadratic equations are often written in the standard form $ax^2 + bx + c = 0$ .
quadratic formula	A formula used to find the solutions of a quadratic equation written in standard form $ax^2 + bx + c = 0$ . The formula is $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$
quadratic term	A term of degree 2. Example: $x^2$ is a quadratic term.

Term	Definition
quadrilateral	A four-sided polygon.
quotient	The result of dividing a number by another number.
radius	A line segment connecting a point on a circle with the center of the circle. The radius is half the length of the diameter.
range of a data set	The difference between the largest number and the smallest number in a set of data.
range of a function	The elements of the second set to which elements of the first set have been assigned.
rate	A comparison of two quantities that have different units of measure. Example: miles per hour, cans per dollar, miles per gallon, and dollars per pound.
ratio	A comparison of two numbers, often expressed by a fraction. Example: if there are four boys in class for every five girls, the ratio of boys to girls is $\frac{4}{5}$ or 4:5 which is read as 4 to 5.
rational numbers	The set of numbers that can be expressed in the form $\frac{a}{b}$ , where $a$ and $b$ are integers and $b$ is not equal to zero. Example: $\frac{6}{5}$ , $-\frac{3}{7}$ , $\frac{4}{1}$
rationalizing the denominator	The process of eliminating a radical from a denominator.
ray	A portion of a line that extends from a given point in one direction only.
real numbers	The set of all rational and irrational numbers.
reciprocal	The reciprocal of a non-zero real number is equal to one divided by that number. The product of a number and its reciprocal is one. Example: The reciprocal of 5 is $\frac{1}{5}$ .
rectangle	A parallelogram with four congruent angles.
rectangular prism	A three-dimensional figure whose sides are all rectangles.
reflection	A transformation that flips a figure across a fixed line, resulting in a mirror image.
reflection symmetry	A two-dimensional figure has reflection symmetry if a line exists about which the figure can be reflected so that the image of the reflection matches the original figure exactly. This line is called the line or axis of symmetry.
regroup	Rearrange a group or groups. Example: 2 tens and 15 ones can be regrouped to 3 tens and 5 ones
regular polygon	A convex polygon with all sides congruent and all angles congruent.
regular pyramid	A pyramid where the base is a regular polygon, and the height meets that base at its center.

<b>Term</b>	<b>Definition</b>
relation	A mathematical relationship between two sets of numbers. A relation may be represented as a set of ordered pair.
relatively prime	Two integers are relatively prime if their greatest common factor is 1. Example: 8 and 15 are relatively prime, but 8 and 12 are not, because their greatest common factor is 4
relevant information	Pertinent or useful information for solving a problem.
remainder	A whole number that is left over after one whole number is divided by another, non-zero whole number.
repeated addition	A way of showing multiplication as a sum of factors. Example: $6 \cdot 5 = 6 + 6 + 6 + 6 + 6$
repeated multiplication	Another way to write numbers in exponential form. Example: $3^4$ can be written as $3 \cdot 3 \cdot 3 \cdot 3$
repeating decimal	A decimal with a block of one or more digits that repeats without end. Example: $\frac{3}{11}$ can be expressed as the repeating decimal $0.\overline{27}$ .
repeating pattern	A sequence whose elements repeat in a consistent way.
rhombus	A parallelogram with four congruent sides.
right angle	An angle whose measure is exactly $90^\circ$ .
right cone	A cone where one end point of the height is at the center of the base and is perpendicular to the base.
right cylinder	A cylinder in which the height from the center of one base meets the center of the other base.
right rectangular prism	A prism with rectangular bases that are perpendicular to its lateral faces.
right triangle	A triangle with one $90^\circ$ angle.
rigid motion	A transformation of the plane or space that preserves distance and angles.
rise	The vertical distance, or change in $y$ , between two points on a coordinate grid.
roots	The solutions of a quadratic equation.
rotation	A transformation that turns a figure around a fixed point.
rotation symmetry	A two-dimensional figure has rotation symmetry if a point exists about which the figure can be rotated less than a full $360^\circ$ turn so that the image of the rotation matches the original figure exactly. This point is called the center of rotation.
round	To arrive at an approximation of a number by following a prescribed algorithm.
rule	A set of operations that corresponds to a pattern or relationship. A rule can be represented as a mathematical expression or equation, or as a written description.
run	The horizontal distance, or change in $x$ , between two points on a coordinate grid.
sample space	The set of all possible outcomes for an experiment.

<b>Term</b>	<b>Definition</b>
scale	An arrangement of numbers that increase by equal intervals. In a graph, there may be a scale for the horizontal axis, for the vertical axis, for both axes, or for neither axis.
scale drawing	A reduced or enlarged drawing whose shape is the same as the actual object, and whose size is determined by a scale factor.
scale factor	The ratio used to compare measurements on a scale drawing to the same measurement on the actual object.
scalene triangle	A triangle that does not have any congruent sides.
scatter plot	A graph of the points representing a collection of data.
scientific calculator	A calculator that represents very large or very small numbers in scientific notation with the powering, factorial, square root, negative and reciprocal keys.
scientific notation	A way of writing very large or very small numbers as a product of a number with an absolute value greater than or equal to 1 and less than 10, and a power of 10. Example: $0.00035 = 3.5 \times 10^{-4}$ or $52,600,000 = 5.26 \times 10^7$
semicircle	An arc of a circle that shares the endpoints of the diameter and is equivalent to half of the circumference.
sequence	Numbers arranged according to a particular pattern.
set	A collection of numbers, objects, or events.
side of a polygon	One of the line segments that enclose a polygon.
Sieve of Eratosthenes	A method of finding all the prime numbers in a certain range. To find all the primes less than 100, start with 2, cross out all numbers greater than 2 that are multiples of 2. Go to the next remaining number, which is 3. Cross out all numbers greater than 3 that are multiples of 3. Go to the next remaining number, which is 5. Cross out all remaining numbers greater than 5 that are multiples of 5, and so on. At each stage, the next number is always a prime. At the end of this process, when there are no more numbers less than 100 to be crossed out, every remaining number is a prime.
similar figures	Figures that have the same shape but not necessarily the same size. Corresponding angles are congruent and the lengths of corresponding side are proportional.
simple event	An event whose probability can be obtained from consideration of a single occurrence. Example: tossing of a coin
simplified algebraic fraction	A fraction in which the numerator and denominator have no common factors other than 1.
simplify an algebraic expression	To combine like terms to write an expression in simplest form.
simplified form of a fraction	A fraction in which the numerator and denominator are relatively prime. Example: $\frac{6}{15} = \frac{2}{5}$
simulation	Modeling a real event without actually observing the event.

Term	Definition
skew lines	Lines that are not in the same plane and are therefore neither parallel nor intersecting.
skip counting	A method of counting that begins with a base integer and then passes over a single integer or set of integers. The pattern for skip counting is often multiples of the base number. Example: skip counting by 2s gives the sequence 2, 4, 6, 8, 10... and skip counting by 5s gives the sequence 5, 10, 15, 20, 25,...
slant height	An altitude of a face of a cone or pyramid.
slope	The steepness of a line graphed on the coordinate plane. It is written as a ratio that compares the change in $y$ values (rise) to the change in $x$ values (run). Lines that rise up from left to right have positive slopes. Lines that fall down from left to right have negative slopes.
slope formula	The formula for finding the slope of a line that passes through points $(x_1, y_1)$ and $(x_2, y_2)$ is $m = \frac{y_2 - y_1}{x_2 - x_1}$ .
slope-intercept form	A linear equation in the form $y = mx + b$ , where $m$ is the slope and $b$ is the $y$ -intercept.
solid figure	A closed three-dimensional figure. Example: cube, sphere, cylinder, pyramid
solution	A value or mathematical representation that satisfies the conditions of a problem.
solution to an equation	Any value of a variable that changes an open sentence into a true statement. Example: 4 is a solution to the equation $x + 6 = 10$ because $4 + 6 = 10$ is a true sentence.
solving	The process of determining the answer to a problem.
solving an equation	Finding the value of the variable for which an equation is true.
square	A parallelogram with four congruent sides and four congruent angles.
square root	One of the two equal factors of a number. Geometrically, the square root is the length of the side of a square that has an area equal to the number for which the square root is being found. Example: $\sqrt{25} = 5$ , because a square with an area of 25 square units has sides measuring 5 units each in length.
square units	The units used to measure area in two dimensions.
squared	Another way to say raised to the second power.
standard form	The way numbers are usually written. Example: $7^2 = 49$ , standard form is 49, and $7^2$ is written in exponential form
standard form of a line	In standard form, a linear function is written as $Ax + By = C$ , where $A$ , $B$ , and $C$ are constants.
statistics	The study of collecting, organizing, and interpreting data.
stem-and-leaf plot	A display of numerical data in a table using digits on the left side as stems and digits on the right side as leaves.

Term	Definition
step plot	A type of graph that represents change in whole increments, represented by a series of discontinuous steps.
straight angle	An angle whose measure is exactly $180^\circ$ .
straightedge	An instrument used to draw straight lines.
subscript	A letter or number used to index a variable. Subscripts are usually set in small type below and to the right of the variable.
substitute	To replace a variable in an equation or expression with a number or expression having equal value.
subtraction	The inverse operation of addition. An operation on two numbers to obtain a third number, called the difference. In the equation $15 - 7 = 8$ , the number 8 is the difference.
subtrahend	In subtraction, the second number in the expression. Example: $15 - 7 = 8$ , the subtrahend is 7.
sum	The result of adding two or more numbers.
sum of the interior angles of a polygon	The total number of degrees in all angles inside of a polygon. The sum of the interior angles of a convex polygon is equal to $180^\circ (n - 2)$ where $n$ is the number of sides of the polygon.
supplementary angles	A pair of angles the sum of whose measures is $180^\circ$ .
surface area	The area of the surface or surfaces of a figure.
symmetric	A shape is symmetric, or symmetrical, if a line can be drawn through the shape such that a reflection across the line creates an identical figure, or if a point can be drawn inside the shape such that a rotation with respect to the point creates an identical figure.
symmetry	The characteristic of a figure that can be divided into congruent sections such that the sections are mirror images of each other.
system of equations	A number of equations with an equal number of variables. The brace indicates a system of equations to be solved simultaneously. Example: $\begin{cases} y = 3x + 2 \\ y = 5x - 8 \end{cases}$
table	A display of information or data by which the facts can be easily read or understood.
tally chart	A chart that uses tally marks to represent data.
tangent to a circle	A line that intersects a circle at exactly one point.
tenth	One of 10 equal parts.
term of a sequence	Any of the numbers in a sequence.
terminating decimal	A decimal that ends. Example: $\frac{1}{4} = 0.25$ which is a terminating decimal.

<b>Term</b>	<b>Definition</b>
terms	The parts of a variable expression that are separated by addition or subtraction signs. These can be numbers, variables, or the product of numbers and variables. Example: In the expression $2x + 5$ , the terms are $2x$ and $5$ .
tessellation	An interlocking pattern of polygons covering a plane with no gaps or overlaps.
theoretical probability	Probability that is expressed as the ratio of the number of times an event can happen to the total number of possible outcomes. This is a probability based on what is likely to occur if an experiment takes place.
transformation	A movement that does not change the shape of a figure.
translation	A transformation of a figure to a new position without turning or flipping the figure.
transversal	A line that intersects two or more coplanar lines.
trapezoid	A four-sided polygon with exactly one pair of parallel sides. The parallel sides are called bases. The two pairs of angles that share a base as a common side are called base angles.
tree diagram	A diagram that shows all of the possible combinations of elements or outcomes of an event.
triangle	A three-sided polygon. The sum of the interior angles of a triangle is $180^\circ$ .
trichotomy property	For all real numbers $a$ and $b$ , exactly one of the following is true: $a < b$ , $a = b$ , or $a > b$ .
trigonometric function	A function (e.g., sine, cosine, tangent, cotangent, secant, cosecant) whose independent variable is an angle measure, usually in degrees or radians.
trigonometric ratio	The ratio of the measures of two sides of a right triangle. There are six basic trigonometric ratios: sine (sin), cosine (cos), tangent (tan), secant (sec), cosecant (csc), and cotangent (cot).
trinomial	A polynomial that consists of three terms, or monomials, when written in standard form.
undefined slope	The slope of a vertical line. The slope is undefined because the difference in the $x$ -values is zero.
unit factor	A special type of ratio that is equivalent to 1. When a number is multiplied by a unit factor, the measurement unit changes, but the value of the measure does not.
unit price	The cost of one item, also called the unit cost.
unlikely	The description of a result that may occur, but probably will not.
variable	An unknown quantity. A variable may be represented by a letter.
variable expression	A mathematical phrase that contains at least one variable.

Term	Definition
velocity	The speed or rate of time at which something moves. The formula for velocity is $v = \frac{d}{t}$ , where $v$ represents velocity, $d$ represents distance, and $t$ represents time.
Venn diagram	A visual device used to represent the relationships among sets.
vertex	The point shared by two rays or line segments forming an angle. It can also refer to the point at which two lines intersect. The plural of vertex is vertices.
vertex of a parabola	The maximum or minimum point of a parabola.
vertex of a polygon	The point at which two sides of a polygon meet.
vertex of an angle	The point where two rays meet to form an angle.
vertical angles	Pairs of opposite angles formed by intersecting lines. Vertical angles are congruent.
vertical axis	The axis running top to bottom, often denoted as the $y$ -axis in a coordinate plane.
vertical line test	A rule used to determine if a graph represents a function. The rule states that if a vertical line cannot be drawn anywhere through the graph so that the line intersects more than one point on the graph, then the graph represents a function.
volume	The number of cubic units inside a three-dimensional figure.
volume formula for a prism or cylinder	The formula $V = Bh$ , where $B$ is the area of the base and $h$ is the height of the prism or cylinder.
volume formula for a pyramid or cone	The formula $V = \frac{1}{3}Bh$ , where $B$ is the area of the base and $h$ is the height of the pyramid or cone.
whole numbers	The set of natural numbers and zero. $\{0, 1, 2, 3, 4, \dots\}$
width	A dimension of a rectangle, usually the shorter side.
$x$ -axis	The horizontal axis in the coordinate plane.
$x$ -intercept	The $x$ -coordinate at which a graph crosses (or intercepts) the $x$ -axis.
$y$ -axis	The vertical axis in the coordinate plane.
$y$ -intercept	The $y$ -coordinate at which a graph crosses (or intercepts) the $y$ -axis. A line can be identified by its slope and $y$ -intercept in slope-intercept form, $y = mx + b$ .
zero exponent	Any term with zero as the exponent is equal to one.
zero property of multiplication	If $a$ and $b$ are real numbers and $ab = 0$ , then $a = 0$ or $b = 0$ .
zero slope	The slope of a horizontal line. In other words, the line does not slant up or down.
zero-product property	If $ab = 0$ , then $a = 0$ or $b = 0$ , and if $a = 0$ or $b = 0$ , then $ab = 0$ .